

Name:	Index No.:	Class:
-------	------------	--------

PRESBYTERIAN HIGH SCHOOL



ADDITIONAL MATHEMATICS Paper 2

4049/02

26 August 2021

Thursday

2 hrs 15 min

*PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL
PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL
PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL
PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL*

2021 SECONDARY FOUR EXPRESS PRELIMINARY EXAMINATIONS

Marking Scheme

This question paper consists of **17** printed pages (including this cover page) and **1** blank page.

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n$$

where n is a positive integer and
$$\binom{n}{r} = \frac{n!}{(n-r)! r!} = \frac{n(n-1)\dots(n-r+1)}{r!}$$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} ab \sin C$$

Answer all questions in the space provided.

1 The equation of a curve is $y = x(4 + x^2)^3$.

- (i) Find an expression for $\frac{dy}{dx}$. [3]

$$\frac{dy}{dx} = x \times 3(4 + x^2)^2 (2x) + (4 + x^2)^3 \times 1$$

M1, M1

$$\frac{dy}{dx} = (4 + x^2)^2 (6x^2 + 4 + x^2)$$

$$\frac{dy}{dx} = (4 + x^2)^2 (7x^2 + 4)$$

A1

- (ii) Explain whether y is an increasing or decreasing function. [2]

For all x, $(4 + x^2)^2 > 0$ and $(7x^2 + 4) > 0$, $\frac{dy}{dx} = (4 + x^2)^2 (7x^2 + 4) > 0$
y is a increasing function. M1
A1

2 The equation of the curve is $y = (a+3)x^2 + ax + 1$, where a is a constant.

(a) When $a = 4$, find the set of values of x for which $y - 4 > 0$. [3]

$$y - 4 > 0$$

$$7x^2 + 4x + 1 - 4 > 0 \quad \text{M1}$$

$$7x^2 + 4x - 3 > 0$$

$$(x+1)(7x-3) > 0 \quad \text{M1}$$

$$x < -1 \text{ or } x > \frac{3}{7} \quad \text{A1}$$

(b) (i) Find the range of values of a for which the curve has no real roots. [3]

$$b^2 - 4ac < 0$$

$$a^2 - 4(a+3)(1) < 0 \quad \text{M1}$$

$$a^2 - 4a - 12 < 0$$

$$(a+2)(a-6) < 0 \quad \text{M1}$$

$$-2 < a < 6 \quad \text{A1}$$

(ii) Hence explain why the curve cannot lie completely below the x -axis. [1]

For curve to lie below x -axis, $a < -3$, but $-2 < a < 6$, hence the curve cannot lie completely below the x -axis

- 3 The expression $f(x) = x^3 + ax + b$, where a and b are constants, is exactly divisible by $x - 2$ and leaves a remainder of 30 when divided by $x - 3$.

- (i) Find the value of a and of b . [4]

$$f(x) = x^3 + ax + b$$

$$\text{when } f(-2) = 0,$$

$$(-2)^3 + a(-2) + b = 0$$

M1

$$-8 - 2a + b = 0 \quad \text{----- (1)}$$

$$\text{when } f(3) = 30,$$

$$(3)^3 + a(3) + b = 30$$

M1

$$27 + 3a + b = 30 \quad \text{----- (2)}$$

$$(2) - (1),$$

$$35 + 5a = 30$$

$$5a = -5$$

$$a = -1$$

A1

$$b = 8 - 2$$

$$b = 6$$

A1

- (ii) Determine by showing all necessary working, the number of real root(s) of the equation $f(x) = 0$. [4]

(ii) $f(x) = x^3 - x + 6$

$$f(x) = (x + 2)(x^2 - 2x + 3)$$

M1, M1

$$\text{Discriminant of } (x^2 - 2x + 3) = (-2)^2 - 4(1)(3) = -8 < 0$$

$$\text{There is no real solution of } (x^2 - 2x + 3)$$

M1

$$f(x) = 0 \text{ has only one real roots}$$

A1

- 4 (i) Prove the identity $\frac{1}{\sec A + 1} + \frac{1}{\sec A - 1} = 2 \operatorname{cosec} A \cot A$. [4]

$$\frac{1}{\sec A + 1} + \frac{1}{\sec A - 1}$$

$$= \frac{\sec A - 1 + \sec A + 1}{\sec^2 A - 1} \quad \text{M1}$$

$$= \frac{2 \sec A}{\tan^2 A} \quad \text{M1}$$

$$= \frac{2 \sec A}{\tan A} \times \frac{1}{\tan A}$$

$$= \frac{2}{\sin A} \times \cot A \quad \text{M1}$$

$$= 2 \operatorname{cosec} A \cot A \quad \text{AG1}$$

- (ii) Hence find all the angles between 0 and 2π for which $\frac{1}{\sec A + 1} + \frac{1}{\sec A - 1} = \frac{4}{3}$. [5]

$$\frac{1}{\sec A + 1} + \frac{1}{\sec A - 1} = \frac{4}{3}$$

$$2 \operatorname{cosec} A \cot A = \frac{4}{3}$$

$$2 \left(\frac{1}{\sin A} \right) \left(\frac{\cos A}{\sin A} \right) = \frac{4}{3} \quad \text{M1}$$

$$\frac{\cos A}{\sin^2 A} = \frac{2}{3}$$

$$3 \cos A = 2 \sin^2 A \quad \text{M1}$$

$$3 \cos A = 2(1 - \cos^2 A) \quad \text{M1}$$

$$3 \cos A = 2 - 2 \cos^2 A$$

$$2 \cos^2 A + 3 \cos A - 2 = 0$$

$$(2 \cos A - 1)(\cos A + 2) = 0$$

$$\cos A = 0.5 \text{ or } \cos A = -2 \text{ (rejected)} \quad \text{M1}$$

$$A = \frac{\pi}{3}, 2\pi - \frac{\pi}{3}$$

$$A = \frac{\pi}{3}, \frac{5\pi}{3}$$

A1

5 It is given that $f(x)$ is such that $f'(x) = \cos 2x + \sin 3x$.

Given that $f(\pi) = 0$, show that $f''(x) + 9f(x) = a + b \sin 2x$ where a and b are constants. [7]

$$f'(x) = \cos 2x + \sin 3x$$

$$f''(x) = -\sin 2x \times 2 + \cos 3x \times 3$$

M1, M1

$$f''(x) = -2 \sin 2x + 3 \cos 3x$$

$$f(x) = \int \cos 2x + \sin 3x \, dx$$

$$f(x) = \frac{\sin 2x}{2} - \frac{\cos 3x}{3} + c$$

M1, M1

$$0 = \frac{\sin 2(\pi)}{2} - \frac{\cos 3(\pi)}{3} + c$$

M1

$$0 = 0 - \frac{-1}{3} + 0 + c$$

$$c = -\frac{1}{3}$$

$$f(x) = \frac{\sin 2x}{2} - \frac{\cos 3x}{3} - \frac{1}{3}$$

$$f''(x) + 9f(x) = -2 \sin 2x + 3 \cos 3x + 9 \left(\frac{\sin 2x}{2} - \frac{\cos 3x}{3} - \frac{1}{3} \right)$$

M1

$$= -2 \sin 2x + 3 \cos 3x + \left(\frac{9 \sin 2x}{2} - 3 \cos 3x - 3 \right)$$

$$= -3 + \frac{5 \sin 2x}{2}$$

A1

- 6 The table below shows the experimental values of two variables x and y .

x	1	2	3	4	5	6
y	63	127	258	510	1000	2100
$\lg y$	1.80	2.10	2.41	2.71	3.00	3.32

It is known that x and y are related by an equation of the form $y = \frac{b^x}{10^a}$, where a and b are constants.

- (i) By plotting $\lg y$ against x , obtain a straight line graph to represent the above data.

[3]

- (ii) Use your graph to estimate the value of a and of b .

[3]

- (ii) Use your graph to find the value of x when $y = 160$.

[1]

- (iii) Explain how would you use the graph to find the value of x for which $(10b)^x = 10^{a+1}$.

[2]

$$(10b)^x = 10^{a+1}$$

$$10^x b^x = 10^a \times 10$$

$$\frac{b^x}{10^a} = \frac{10}{10^x}$$


M1

$$y = 10^{1-x}$$

$$\lg y = \lg(10^{1-x})$$

$$\lg y = 1 - x$$

Draw the line $\lg y = 1 - x$ and find the x -coordinate of the point of intersection. A1



- 7 (i) Express $\frac{5x^2 - 12x - 3}{(x-2)(x^2+3)}$ in partial fractions. [5]

$$\frac{5x^2 - 12x - 3}{(x-2)(x^2+3)} = \frac{A}{x-2} + \frac{Bx+C}{x^2+3}$$

M1

$$5x^2 - 12x - 3 = A(x^2 + 3) + (Bx + C)(x - 2)$$

Let $x = 2$

$$5(2)^2 - 12(2) - 3 = A(2^2 + 3)$$

M1 for at least one correct method

$$-7 = 7A$$

$$-1 = A$$

Let $x = 0$

$$-3 = -1(0^2 + 3) + (C)(0 - 2)$$

$$-3 = -3 - 2C$$

$$0 = C$$

A1 for at least one correct
A2 for all 3 correct

Let $x = 1$

$$5(1)^2 - 12 - 3 = -1(1^2 + 3) + (B)(1 - 2)$$

$$-10 = -4 - B$$

$$6 = B$$

$$\frac{5x^2 - 12x - 3}{(x-2)(x^2+3)} = -\frac{1}{(x-2)} + \frac{6x}{(x^2+3)}$$

A1

- (ii) Differentiate $\ln(x^2 + 3)$ with respect to x . [1]

$$\begin{aligned}\frac{d[\ln(x^2 + 3)]}{dx} &= \frac{1}{x^2 + 3} \times 2x \\ &= \frac{2x}{x^2 + 3}\end{aligned}$$

B1

- (iii) Using the results from (i) and (ii), determine $\int \frac{5x^2 - 12x - 3}{(x-2)(x^2 + 3)} dx$. [4]

$$\int \frac{5x^2 - 12x - 3}{(x-2)(x^2 + 3)} dx = \int -\frac{1}{x-2} + \frac{6x}{x^2 + 3} dx$$

M1


$$\int \frac{5x^2 - 12x - 3}{(x-2)(x^2 + 3)} dx = \int -\frac{1}{x-2} dx + 3 \int \frac{2x}{x^2 + 3} dx$$

M1

$$\int \frac{5x^2 - 12x - 3}{(x-2)(x^2 + 3)} dx = -\ln(x-2) + 3\ln(x^2 + 3) + c$$

A1, A1

- 8 The diagram shows a rectangle $ABCD$. A line through A intersects CD at F and BC produced at E . It is given that $\angle BAE = \theta^\circ$, $AF = 18$ cm and $FE = 16$ cm.



- (i) Show that the perimeter, P cm, of the rectangle is given by

$$P = 68 \cos \theta + 36 \sin \theta.$$

[3]

In $\triangle AFD$, $\cos \theta = \frac{DF}{18}$

$\cos \theta$ and
 $\sin \theta$ of
either
triangle

$$DE = 18 \cos \theta$$

$$\sin \theta = \frac{AF}{18}$$

M1

$$AF = 18 \sin \theta$$

In $\triangle FCE$, $\cos \theta = \frac{FC}{16}$

$$FC = 16 \cos \theta$$

M1

$$\sin \theta = \frac{CE}{16}$$

$$CE = 16 \sin \theta$$

Therefore $P = 2(AB + AD)$

$$P = 2(18 \cos \theta + 16 \cos \theta + 18 \sin \theta)$$

$$P = 68 \cos \theta + 36 \sin \theta \text{ (shown)}$$

} AG1

(ii) Express P in the form $R \cos(\theta - \alpha)$. [3]

$$R = \sqrt{68^2 + 36^2} = \sqrt{5920} \quad \text{M1}$$

$$\tan \alpha = \frac{36}{68} \quad \text{M1}$$

$$\alpha = 27.9^\circ$$

$$P = \sqrt{5920} \cos(\theta - 27.9^\circ) \quad \text{A1}$$

(iii) Find the value of θ when the perimeter is 62 cm.

[2]

$$\sqrt{5920} \cos(\theta - 27.9^\circ) = 62 \quad \text{M1}$$

$$\cos(\theta - 27.9^\circ) = \frac{62}{\sqrt{5920}}$$

$$\theta - 27.89^\circ = 36.31$$

$$\theta = 64.2^\circ$$

A1

9 A circle C_1 has equation $x^2 + y^2 - 6x + 4y = 12$.

- (i) Find the radius and the coordinates of the centre of C_1 . [3]

$$\text{Centre} = \left(\frac{-6}{-2}, \frac{4}{-2} \right)$$

$$\text{Centre} = (3, -2) \quad \text{B1}$$

$$\text{Radius} = \sqrt{3^2 + (-2)^2 - (-12)} \quad \text{M1 for attempt to find centre or radius}$$

$$\text{Radius} = 5 \quad \text{A1}$$

- (ii) Show that the equation of the tangent to the circle at the point $(7, -5)$ is $3y = 4x + k$ where k is an integer. [3]

$$\text{Gradient of the line from the point of tangent to centre} = \frac{-5 - (-2)}{7 - 3} \quad \text{M1}$$

$$\text{Gradient of the tangent} = -\frac{1}{\left(\frac{3}{-4} \right)} = \frac{4}{3} \quad \text{M1}$$

$$y - (-5) = \frac{4}{3}(x - 7)$$

$$3y + 15 = 4x - 28$$

$$3y = 4x - 43 \quad \text{A1}$$

- (iii) Another circle C_2 has centre $(-8, 4)$ and radius 7 cm. Find the shortest distance between the 2 circles. [2]

$$\text{Distance between the 2 centres} = \sqrt{(3 - (-8))^2 + (-2 - 4)^2} = \sqrt{157} \quad \text{M1}$$

$$\text{Shortest distance} = \sqrt{157} - 5 - 7 = 0.530 \quad \text{A1}$$

- 10 (a) Without using a calculator, find the value of 6^x given that $12^{x-2} = 3^{4-x}$. [4]

$$12^{x-2} = 3^{4-x}$$

$$12^x \times 12^{-2} = 3^4 \times 3^{-x} \quad \text{M1}$$

$$\frac{12^x}{3^{-x}} = \frac{3^4}{12^{-2}} \quad \text{M1}$$

$$36^x = 9^2 \times 12^2$$

$$(6^x)^2 = (9 \times 12)^2 \quad \text{M1}$$

$$6^x = 108 \quad \text{A1}$$

- (b) Solve the equation $\log_2 x = 6 + \log_{16} x$. [3]

$$\log_2 x = 6 + \log_{16} x$$

$$\log_2 x = 6 + \frac{\log_2 x}{\log_2 16} \quad \text{M1 (Change of base)}$$

$$\log_2 x = 6 + \frac{\log_2 x}{4}$$

$$\frac{3 \log_2 x}{4} = 6 \quad \text{M1}$$

$$\log_2 x = 8$$

$$x = 2^8$$

$$x = 256 \quad \text{A1}$$

(c) Solve the equation $\lg(x-4) + 2\lg 3 = 1 + \lg\left(\frac{x}{2}\right)$. [3]

$$\lg(x-4) + 2\lg 3 = 1 + \lg\left(\frac{x}{2}\right)$$

$$\lg(x-4) + \lg 3^2 = \lg 10 + \lg\left(\frac{x}{2}\right)$$

M1 (Power rule)

$$\lg 9(x-4) = \lg 10\left(\frac{x}{2}\right)$$


M1 (Product rule)

$$9x - 36 = 5x$$

$$4x = 36$$

$$x = 9$$

A1



The diagram shows part of the curve $y = 27 + \frac{8}{(x-2)^3}$ which crosses the axes at points A and B

- (i) Find the coordinates of A and of B. [2]

$$\text{When } x = 0, y = 27 + \frac{8}{(0-2)^3} = 26$$

$$B(0, 26)$$

B1

$$\text{When } y = 0,$$

$$0 = 27 + \frac{8}{(x-2)^3}$$

$$-27 = \frac{8}{(x-2)^3}$$

$$(x-2)^3 = \frac{8}{-27}$$

$$x-2 = -\frac{2}{3}$$

$$x = \frac{4}{3}$$

$$\left(\frac{4}{3}, 0\right)$$

B1

The normal to the curve at B cuts the x -axis at C .

Find

- (ii) the equation of the normal, [3]

$$\frac{dy}{dx} = -\frac{24}{(x-2)^4} \quad \text{M1}$$

$$\frac{dy}{dx} = -\frac{24}{(0-2)^4} = -\frac{3}{2} \quad \text{M1}$$

$$\text{Gradient of normal} = \frac{2}{3}$$

$$\text{Equation of normal is } y = \frac{2}{3}x + 26 \quad \text{A1}$$

- (iii) the area of the shaded region. [4]

$$0 = \frac{2}{3}x + 26$$

$$-\frac{2}{3}x = 26$$

$$x = -39 \quad \text{M1}$$

$$\begin{aligned} \text{Area of shaded region} &= \frac{1}{2}(39)(26) + \int_0^4 27 + 8(x-2)^{-3} dx \\ &= 507 + \left[27x + \frac{8}{-2}(x-2)^{-2} \right]_0^4 \end{aligned} \quad \begin{array}{l} \text{M1 for} \\ \text{either} \\ \text{M1 for} \\ \text{integratio} \\ \text{n} \end{array}$$

$$= 507 + \left[27\left(\frac{4}{3}\right) - 4\left(\frac{4}{3} - 2\right)^{-2} - \left(27(0) - 4(0-2)^{-2}\right) \right]$$

$$= 507 + [36 - 9 + 1]$$

$$= 535 \text{ unit}^2 \quad \text{A1}$$

END OF PAPER