




CANBERRA SECONDARY SCHOOL



## 2021 Preliminary Examination

### Secondary Four Express

#### ADDITIONAL MATHEMATICS

4049/02

27<sup>th</sup> August 2021  
2 hours 15 minutes  
0800h – 1015h

Name: \_\_\_\_\_ (     )     Class: \_\_\_\_\_

#### READ THESE INSTRUCTIONS FIRST

Write your full name, class and index number on all work you hand in.  
Write in dark blue or black pen on both sides of the paper.  
You may use a pencil for any diagrams or graphs.  
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.

If working is needed for any question it must be shown with the answer.

Omission of essential working will result in loss of marks.

Calculators should be used where appropriate.

If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place.

For  $\pi$ , use either your calculator value or 3.142, unless the question requires the answer in terms of  $\pi$ .

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total number of marks for this paper is 90.

FOR MARKER'S USE		
	Marks Awarded	Max Marks
Total		90

---

This question paper consists of 19 printed pages including the cover page.

Setter: Mr Lathif and Mrs Wee

**Mathematical Formulae****1. ALGEBRA***Quadratic Equation*

For the equation  $ax^2 + bx + c = 0$ ,  $a \neq 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

*Binomial expansion*

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n,$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1)\dots(n-r+1)}{r!}$ .

**2. TRIGONOMETRY***Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

*Formulae for  $\triangle ABC$* 

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

Answer **all** the questions

**1** Given that  $f(x) = 2x^3 + px^2 + 7x + q$ .

Find the value of  $p$  and of  $q$  for which  $(x-3)$  and  $(x-2)$  are factors of  $f(x)$ .


Hence solve  $f(x) = 0$ .

[8]

2 (i) Prove the identity  $\frac{1+\sin x}{\cos x} + \frac{\cos x}{1+\sin x} = 2 \sec x$ . [4]

(ii) Hence, solve the equation  $\frac{1+\sin x}{\cos x} + \frac{\cos x}{1+\sin x} = 4$  for  $0 < x < 2\pi$ . [4]

- 3 The diagram shows part of a straight line obtained by plotting  $x^2y$  against  $x$ .




- (i) Find an expression for  $y$  in terms of  $x$ .

[4]

(ii) Find  $x$  when  $y = \frac{5}{x^2}$ . [2]

(iii) For the above equation found in part (i), suggest another expression for the vertical and horizontal axes such that a straight line can also be plotted. What is the value of the gradient and  $y$  – intercept of this new equation? [3]

- 4 The diagram below shows the side view of a warehouse  $ABCD$  such that  $BC = 10$  m,  $CD = 6$  m,  $\angle BCD = \angle BAD = 90^\circ$  and  $\angle CBA = \theta$ .



- (i) Show that  $AB = 6 \sin \theta + 10 \cos \theta$ .

[2]

(ii) Express  $AB$  in the form  $R \sin(\theta + \alpha)$ , where  $R$  is positive and  $\alpha$  is an acute angle.

Hence, find the maximum value of  $AB$  and the value of  $\theta$  for which this occurs.

[6]

(iii) Explain why this value of  $\theta$  is the smallest possible.

[2]

5 Points  $A(3,8)$  and  $B(6,5)$  are on a circle.

(i) Find the equation of perpendicular bisector of  $AB$ . [3]

(ii) Given that  $y + x = 8$  passes through the center of the circle, find the coordinates of the center and the radius of the circle. [3]

(iii) Hence, find the equation of the circle. [1]

(iv) Explain if point  $C(6,3)$  is inside or outside the circle. [2]

(v) Which axis is tangent to the circle? Explain your answer. [2]

6 (a) Given that  $y = \sec x$ , show that  $\frac{dy}{dx} = \sec x \tan x$  .

Hence show that  $y^4 - y^2 = \left(\frac{dy}{dx}\right)^2$  .

[5]


- (b) Given that  $\frac{6x+1}{2x+1}$  can be written in the form  $A + \frac{B}{2x+1}$ , find  $A$  and  $B$ .

Hence, or otherwise  $\int \frac{6x+1}{2x+1} dx$ .

[5]

- 7 A cone of height,  $h$  and radius,  $r$  fits exactly inside a sphere of radius 5 cm as shown below.

$$\left( \text{Volume of cone} = \frac{1}{3} \pi r^2 h \right)$$




- (i) Express  $r$  in terms of  $h$ . [2]

- (ii) Show that the volume of cone is given by  $V = \frac{\pi h^2 (10 - h)}{3}$ . [2]

(iii) Find the value of  $h$  for which  $V$  is stationary. [3]

(iv) Find the volume of the cone for this value of  $h$  and determine if this volume is a maximum or minimum.  
Justify your answer. [4]

- 8 The diagram below shows the curve  $y = x^3 - 12x^2 + 20x$ , which crosses the  $x$ -axis at the origin  $O$  and the points  $A$  and  $B$ .



- (i) Find the coordinates of  $A$  and  $B$ .

[3]

The tangent to the curve at the point  $(8, -96)$  crosses the  $x$  – axis at the point  $C$ .

**(ii)** Find the equation of the tangent and hence, the coordinates of  $C$ . [4]

**(iii)** Find the area of the shaded region. [4]

**9 (a)** Solve  $9^{x+1} - 3^{x+2} = 2(3^{x+1} - 2)$ . [4]

**(b)** Given that  $\log_3(2x-1) = \log_9(2x-1) + 1$ , find the value of  $x$ . [3]

- (c) Given that  $\log_a xy = m$  and  $\log_a \frac{x^3}{y^2} = n$ , find  $5\log_a \left(\frac{x}{y}\right)$  in terms of  $m$  and  $n$ .

[5]

<b>1</b>	$2x^3 - 9x^2 + 7x + 6 = 0$ $(2x+1)(x-3)(x-2) = 0$ $x = -\frac{1}{2}, x = 3 \text{ or } x = 2$
<b>2ii</b>	$x = \frac{\pi}{3}, \frac{5\pi}{3}$
<b>3i</b>	$y = \frac{2}{3x} + \frac{3}{x^2}$
<b>3ii</b>	$x = 3$
<b>3iii</b>	Plot $xy$ against $\frac{1}{x}$ where the gradient is 3 and the $y$ -intercept is $\frac{2}{3}$ , o.e
<b>4ii</b>	$6\sin\theta + 10\cos\theta = \sqrt{136}\sin(\theta + 59.0^\circ)$ <p>Maximum <math>AB = \sqrt{136}</math> or <math>2\sqrt{34}</math> or 11.7 m when <math>\theta = 31.0^\circ</math></p>
<b>4iii</b>	For $AB = \sqrt{136} = \sqrt{10^2 + 6^2}$ , $AD$ must be 0, so that triangle $BCD$ becomes a right-angled triangle. This will result that $\angle CBA = \theta$ to be smallest possible.
<b>5i</b>	$y = x + 2$
<b>5ii</b>	Center (3,5) Radius = 3 units
<b>5iii</b>	$(x-3)^2 + (y-5)^2 = 9$
<b>5iv</b>	C is outside of the circle.
<b>5v</b>	Since the $x$ -coordinate of the center of the circle is equal to the radius, the $y$ -axis is tangent to the circle.
<b>6b</b>	$3x - \ln(2x+1) + c$
<b>7i</b>	$r = \sqrt{10h - h^2}$
<b>7iii</b>	$h = 6\frac{2}{3} \text{ cm}$
<b>7iv</b>	$V = 155 \text{ cm}^3$ (3 s.f.) Area is maximum.
<b>8i</b>	$A(2,0)$ and $B(10,0)$
<b>8ii</b>	$C(12.8,0)$
<b>8iii</b>	$638.4 \text{ units}^2$
<b>9a</b>	$x = -1$ or $x \approx 0.262$
<b>9b</b>	$x = 5$
<b>9c</b>	$5\log_a\left(\frac{x}{y}\right) = 2n - m$