

Name:	Class:	Class Register Number:
-------	--------	------------------------



**中正中学**

**CHUNG CHENG HIGH SCHOOL (MAIN)**

Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School  
 Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School  
 Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School  
 Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School  
 Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School

**PRELIMINARY EXAMINATION 2023  
SECONDARY 4**

**ADDITIONAL MATHEMATICS**

**4049/01**

**Paper 1**

**Thursday 24 August 2023**

**2 hours 15 minutes**

**MARKS SCHEME**

This document consists of **19** printed pages and **1** blank page.

**[Turn over**

**Mathematical Formulae****1. ALGEBRA***Quadratic Equation*

For the equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

*Binomial expansion*

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n,$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1) \dots (n-r+1)}{r!}.$

**2. TRIGONOMETRY***Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

*Formulae for  $\Delta ABC$* 

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} ab \sin C$$

- 1 The points  $R$  and  $S$  have coordinates  $(\sqrt{3}, 2\sqrt{3})$  and  $(\sqrt{5}, 4\sqrt{5})$  respectively. Show that the gradient of  $RS$  can be expressed in the form  $a + b\sqrt{15}$ , where  $a$  and  $b$  are integers to be found. [4]

$$\begin{aligned}
 \text{Gradient of } RS &= \frac{4\sqrt{5} - 2\sqrt{3}}{\sqrt{5} - \sqrt{3}} \times \frac{\sqrt{5} + \sqrt{3}}{\sqrt{5} + \sqrt{3}} \\
 &= \frac{(4\sqrt{5} - 2\sqrt{3})(\sqrt{5} + \sqrt{3})}{(\sqrt{5})^2 - (\sqrt{3})^2} \\
 &= \frac{20 - 4\sqrt{15} - 2\sqrt{15} - 6}{2} \\
 &= \frac{14 - 6\sqrt{15}}{2} \\
 &= \frac{2(7 - 3\sqrt{15})}{2} \\
 &= 7 - 3\sqrt{15} \\
 a &= 7, b = -3
 \end{aligned}$$

$$\text{B1} - \frac{4\sqrt{5} - 2\sqrt{3}}{\sqrt{5} - \sqrt{3}}$$

$$\text{M1} - \sqrt{\frac{\sqrt{5} + \sqrt{3}}{\sqrt{5} + \sqrt{3}}}$$

M1 – either numerator or denominator expanded correctly

A1

2 Given that  $\cos \theta = p$  and that  $\theta$  is acute, express in terms of  $p$ ,

(a)  $\sin \theta$ ,

[1]

$$\sqrt{1-p^2}$$

B1

(b)  $\tan(90^\circ - \theta)$ ,

[2]

$$\begin{aligned} &= \frac{1}{\tan \theta} \\ &= \frac{1}{\frac{\sqrt{1-p^2}}{p}} \\ &= \frac{p}{\sqrt{1-p^2}} \end{aligned}$$

$$\text{M1} - \frac{1}{\text{their } \tan \theta}$$

A1

(c)  $\cos 2\theta$ .

[2]

$$\begin{aligned} \cos 2\theta &= 1 - 2\sin^2 \theta \\ &= 1 - 2(\sqrt{1-p^2})^2 \\ &= 1 - 2(1-p^2) \\ &= 1 - 2 + 2p^2 \\ &= 2p^2 - 1 \end{aligned}$$

M1 – uses any  $\cos 2\theta$  formula correctly

A1

3 Express  $\frac{12x^2 + 32x + 31}{(2x-1)(x+2)^2}$  in partial fractions.

[5]

$$\frac{12x^2 + 32x + 31}{(2x-1)(x+2)^2} = \frac{A}{2x-1} + \frac{B}{x+2} + \frac{C}{(x+2)^2}$$

$$12x^2 + 32x + 31 = A(x+2)^2 + B(2x-1)(x+2) + C(2x-1)$$

$$\text{let } x = \frac{1}{2},$$

$$50 = \frac{25}{4}A$$

$$A = 8$$

$$\text{let } x = -2,$$

$$15 = -5C$$

$$C = -3$$

$$\text{let } x = 0,$$

$$31 = 4A - 2B - C$$

$$31 = 32 - 2B + 3$$

$$2B = 4$$

$$B = 2$$

$$A = 8, B = 2, C = -3$$

$$\frac{12x^2 + 32x + 31}{(2x-1)(x+2)^2} = \frac{8}{2x-1} + \frac{2}{x+2} - \frac{3}{(x+2)^2}$$

M1 – realising the form of partial fractions

M1 – realising the need to eliminate the denominator

A1, A1, A1 – do not award last A1 if not expressed in partial fractions

- 4 The expression  $ax^3 + bx^2 + b$  leaves a remainder of  $R$  when divided by  $(x+1)$  and a remainder of  $5R-2$  when divided by  $(x+2)$ .

(a) Show that  $b = \frac{2-3a}{5}$ . [4]

when  $x = -1$ ,

$$a(-1)^3 + b(-1)^2 + b = R$$

$$-a + 2b = R \quad \text{--- (1)}$$

when  $x = -2$ ,

$$a(-2)^3 + b(-2)^2 + b = 5R - 2$$

$$-8a + 5b = 5R - 2 \quad \text{--- (2)}$$

sub (1) into (2),

$$-8a + 5b = 5(-a + 2b) - 2$$

$$-8a + 5b = -5a + 10b - 2$$

$$10b - 5b = -8a + 5a + 2$$

$$5b = -3a + 2$$

$$b = \frac{2-3a}{5} \quad \text{(shown)}$$

M1 – realises  $f(-1) = R$

M1 – realises  $f(-2) = 5R - 2$

DM1 – award only if both M1 above is achieved

A1

AG

- (b) Given further that  $ab = -8$  and  $a > b$ , find the value of  $a$  and of  $b$ . [3]

$$ab = -8$$

$$b = \frac{-8}{a}$$

$$\frac{-8}{a} = \frac{-3a+2}{5}$$

$$-40 = -3a^2 + 2a$$

$$3a^2 - 2a - 40 = 0$$

$$(3a+10)(a-4) = 0$$

$$a = -\frac{10}{3} \quad \text{or} \quad a = 4$$

$$b = 2.4 \quad b = -2$$

(reject)

$$\therefore a = 4, b = -2$$

M1 – allow slips only for LHS

√M1 – finding ‘b’

A1

- (c) Using the values of  $a$  and  $b$  found in **part (b)**, explain why the equation  $ax^3 + bx^2 + b = 0$  has only one real root and state its value. [4]

$$4x^3 - 2x^2 - 2 = 0$$

$$2x^3 - x^2 - 1 = 0$$

when  $x = 1$ ,

$$2(1)^3 - 1^2 - 1 = 0$$

$\therefore$  By factor theorem,  $(x-1)$  is a factor.

$$(x-1)(2x^2 + cx + 1) = 0$$

by comparing  $x^2$  terms,

$$-2 + c = -1$$

$$c = 1$$

$$(x-1)(2x^2 + x + 1) = 0$$

For  $2x^2 + x + 1 = 0$ ,

$$\text{discriminant} = 1^2 - 4(2)(1) = -7 < 0$$

$\therefore 2x^2 + x + 1 = 0$  has no real roots.

$4x^3 - 2x^2 - 2 = 0$  has only 1 real root of  $x = 1$ .

B1 - factor

M1 – correct method to find quadratic (division or inspection)

M1 – realising the need to find discriminant / solve quadratic equation

A1 – no real roots +  $x = 1$

- 5 Find the coordinates of the stationary points of the curve  $y = \frac{(x-3)^2}{x}$  and determine the nature of each stationary point. [7]

$$y = \frac{(x-3)^2}{x}$$

$$= \frac{x^2 - 6x + 9}{x}$$

$$= x - 6 + \frac{9}{x}$$

$$= x - 6 + 9x^{-1}$$

$$\frac{dy}{dx} = 1 - 9x^{-2}$$

$$= 1 - \frac{9}{x^2}$$

For stationary points,  $\frac{dy}{dx} = 0$

$$1 - \frac{9}{x^2} = 0$$

$$\frac{9}{x^2} = 1$$

$$x^2 = 9$$

$$x = 3 \text{ or } -3$$

$$y = 0 \text{ or } -12$$

$$\frac{d^2y}{dx^2} = 18x^{-3}$$

$$= \frac{18}{x^3}$$

At (3,0),

$$\frac{d^2y}{dx^2} = \frac{18}{3^3} = \frac{2}{3} > 0$$

$\therefore (3,0)$  is a minimum point

At (-3,-12),

$$\frac{d^2y}{dx^2} = \frac{18}{(-3)^3} = -\frac{2}{3} < 0$$

$\therefore (-3,-12)$  is a maximum point

B1

M1 – sets  $\frac{dy}{dx}$  to 0

A1 – for (3,0)

A1 – for (-3, -12)

M1 – their 2<sup>nd</sup> derivative or use of 1<sup>st</sup> derivative test

A1

A1



- 6** The height above ground,  $h$  metres, of a ball, released by a machine can be modelled by the equation  $h = -0.2x^2 + 6x + 3$  where  $x$  is the horizontal distance travelled by the ball in metres.

- (a) State the height above ground at which the ball is released. [1]

3 metres

- (b) Express  $h = -0.2x^2 + 6x + 3$  in the form  $a(x-b)^2 + c$ , where  $a$ ,  $b$  and  $c$  are constants to be found. [2]

$$\begin{aligned} h &= -0.2(x^2 - 30x) + 3 \\ &= -0.2(x^2 - 30x + 15^2 - 15^2) + 3 \\ &= -0.2[(x-15)^2 - 225] + 3 \\ &= -0.2(x-15)^2 + 48 \end{aligned}$$

B2, 1 : -1 for each error

- (c) Using your result from (b), explain why the height of the ball can never be more than 48 metres. [2]

$$\text{For } -0.2(x-15)^2 + 48,$$

$$(x-15)^2 \geq 0,$$

$$-0.2(x-15)^2 \leq 0$$

$$-0.2(x-15)^2 + 48 \leq 48$$

Since the ball reaches a maximum height of 48 m, it will never reach a height of more than 48 m.

✓ M1 – their square term

A1

- (d) Hence, explain if this machine is safe for use in an indoor stadium with a ceiling height of 45 metres. [1]

Since the ball can reach a maximum height of 48m which exceeds the ceiling height of 45 m, this machine is not safe for use in the indoor stadium.

B1 – to have comparison of maximum height with ceiling height.

7 (a) Solve the equation  $3^x(18+3^x)=40$ .

[3]

$$3^x(18+3^x)=40$$

$$(3^x)^2+18(3^x)-40=0$$

Let  $y=3^x$ .

$$y^2+18y-40=0$$

$$(y+20)(y-2)=0$$

$$y=-20 \quad \text{or} \quad y=2$$

$$3^x=2$$

$$\ln 3^x=\ln 2$$

$$3^x=-20 \quad (\text{rej.} \because 3^x > 0) \quad \text{or} \quad x \ln 3 = \ln 2$$

$$x = \frac{\ln 2}{\ln 3}$$

$$=0.631 \quad (3 \text{ sig. fig.})$$

M1 – solving of quadratic equation

M1 – taking  $\ln$ 

A1

(b) Solve the equation  $\log_{\sqrt{2}} y = 3 + \log_2(y+6)$ .

[5]

$$\log_{\sqrt{2}} y = 3 + \log_2(y+6)$$

$$\log_{\sqrt{2}} y - \log_2(y+6) = 3$$

$$\frac{\log_2 y}{\log_2 \sqrt{2}} - \log_2(y+6) = 3$$

$$\frac{\log_2 y}{\frac{1}{2}} - \log_2(y+6) = 3$$

$$2\log_2 y - \log_2(y+6) = 3$$

$$\log_2 \left( \frac{y^2}{y+6} \right) = 3$$

Comparing,

$$\frac{y^2}{y+6} = 3$$

$$y^2 = 3y + 18$$

$$y^2 - 3y - 18 = 0$$

$$(y-6)(y+3) = 0$$

$$y = 6 \quad \text{or} \quad -3 \quad (\text{rej.} \because y > 0)$$

B1 – change of base law

M1 – power law

M1 – quotient law

M1 – removal of log/comparing

A1

Continuation of working space for question 7(b)

- (c) In order to obtain a graphical solution of the equation  $x = 2 \ln \left( 4 - \frac{3x}{2} \right)$ , a suitable straight line can be drawn on the same set of axes as the graph of  $y = 4 - e^{\frac{x}{2}}$ . Make  $e^{\frac{x}{2}}$  the subject of  $x = 2 \ln \left( 4 - \frac{3x}{2} \right)$  and hence find the equation of this line. [3]

$$x = 2 \ln \left( 4 - \frac{3x}{2} \right)$$

$$\frac{x}{2} = \ln \left( 4 - \frac{3x}{2} \right)$$

$$e^{\frac{x}{2}} = 4 - \frac{3x}{2}$$

$$\frac{3x}{2} = 4 - e^{\frac{x}{2}}$$

$$\text{Equation of line: } y = \frac{3x}{2}$$

M1 – attempt to make  $\ln \left( 4 - \frac{3x}{2} \right)$  the subject

A1

B1

8 A curve has the equation  $y = \frac{x^2 - x + 1}{5x - 5}$ ,  $x \neq 1$ .

(a) Show that  $\frac{dy}{dx} = \frac{x^2 - 2x}{5(x-1)^2}$ . [3]

$$\begin{aligned}\frac{dy}{dx} &= \frac{(5x-5)(2x-1) - (x^2 - x + 1)(5)}{(5x-5)^2} \\ &= \frac{10x^2 - 5x - 10x + 5 - 5x^2 + 5x - 5}{[5(x-1)]^2} \\ &= \frac{5x^2 - 10x}{5^2(x-1)^2} \\ &= \frac{5(x^2 - 2x)}{25(x-1)^2} \\ &= \frac{x^2 - 2x}{5(x-1)^2} \text{ (shown)}\end{aligned}$$

M1 – quotient rule performed correctly, allow for numerical slips

B1 – for  $5^2(x-1)^2 / 25(x-1)^2$

A1 – factorisation must be seen

AG

(b) Explain why the curve is increasing for  $x > 2$ . [3]

**Alternative solution**

Where the curve is increasing,

$$\frac{dy}{dx} > 0,$$

$$\frac{x^2 - 2x}{5(x-1)^2} > 0$$

Since  $5(x-1)^2 > 0$ , for all real

values of  $x$ ,

$$x^2 - 2x > 0$$

$$x(x-2) > 0$$

$$x < 0 \text{ or } x > 2$$

$\therefore$  for  $x > 2$ , the curve is always increasing.

$$x^2 - 2x = x(x-2)$$

$$x > 2, x-2 > 0$$

$$x(x-2) > 0$$

$$\text{since } (x-1)^2 > 0$$

$$\frac{x^2 - 2x}{5(x-1)^2} > 0$$

Since  $\frac{dy}{dx} > 0$ , the curve is always increasing.

B1 –  $x > 2$ ,  $x-2 > 0$ ,  $x(x-2) > 0$

B1 –  $(x-1)^2 > 0$

B1 –  $\frac{x^2 - 2x}{5(x-1)^2} > 0$  + conclusion of

curve is always increasing. Award only if B1, B1 has been achieved.

- 9 (a) Given that  $\frac{\cos(\alpha + \beta)}{\cos(\alpha - \beta)} = \frac{3}{4}$ , prove that  $\cos \alpha \cos \beta = 7 \sin \alpha \sin \beta$ . [2]

$$\begin{aligned}
 4[\cos \alpha \cos \beta - \sin \alpha \sin \beta] &= 3[\cos \alpha \cos \beta + \sin \alpha \sin \beta] \\
 4 \cos \alpha \cos \beta - 4 \sin \alpha \sin \beta &= 3 \cos \alpha \cos \beta + 3 \sin \alpha \sin \beta \\
 4 \cos \alpha \cos \beta - 3 \cos \alpha \cos \beta &= 4 \sin \alpha \sin \beta + 3 \sin \alpha \sin \beta \\
 \cos \alpha \cos \beta &= 7 \sin \alpha \sin \beta \text{ (shown)}
 \end{aligned}$$

M1 – attempt at addition formula

A1

AG

- (b) Hence, deduce the relationship between  $\tan \alpha$  and  $\tan \beta$ . [2]

$$\begin{aligned}
 \cos \alpha \cos \beta &= 7 \sin \alpha \sin \beta \\
 \frac{1}{7} &= \frac{\sin \alpha \sin \beta}{\cos \alpha \cos \beta} \\
 \tan \alpha \tan \beta &= \frac{1}{7} \quad \text{or} \quad \frac{1}{\tan \beta} = 7 \tan \alpha
 \end{aligned}$$

M1 – realises the need to divide

A1

- (c) Given further that  $\alpha + \beta = 45^\circ$ , calculate the value of  $\tan \alpha + \tan \beta$ . [3]

$$\begin{aligned}
 \tan(\alpha + \beta) &= \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta} \\
 \tan 45^\circ &= \frac{\tan \alpha + \tan \beta}{1 - \frac{1}{7}} \\
 1 &= \frac{\tan \alpha + \tan \beta}{\frac{6}{7}} \\
 \tan \alpha + \tan \beta &= \frac{6}{7}
 \end{aligned}$$

M1 –  $\tan 45^\circ$  or realises the need for  $\frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$

M1- their  $\tan \alpha \tan \beta$

A1

**10** A circle, whose equation is  $x^2 + y^2 - 10x - 8y + 16 = 0$ , has centre  $C$  and radius  $r$ .

**(a)** Find the coordinates of  $C$  and the value of  $r$ .

[3]

$$-2a = -10$$

$$a = 5$$

$$-2b = -8$$

$$b = 4$$

$$a^2 + b^2 - r^2 = 16$$

$$r^2 = 5^2 + 4^2 - 16$$

$$r = 5$$

$\therefore C(5, 4)$  and  $r$  is 5 units.

M1 – their  $a$  and  $b$

B1 – (5,4), A1 – 5 units

### Alternative Solution

$$x^2 + y^2 - 10x - 8y + 16 = 0$$

$$x^2 - 10x + 5^2 - 5^2 + y^2 - 8y + 4^2 - 4^2 + 16 = 0$$

$$(x-5)^2 + (y-4)^2 = 25 + 16 - 16$$

$$(x-5)^2 + (y-4)^2 = 5^2$$

$C(5, 4)$  and  $r = 5$  units.

M1 – attempt to complete square

B1 – (5,4), A1 – 5 units

**(b)** Explain whether the point  $(9, 2)$  lies inside or outside the circle.

[2]

Distance from  $(9, 2)$  to  $C$

$$= \sqrt{(9-5)^2 + (4-2)^2}$$

$$= \sqrt{20} < 5 \text{ (radius)}$$

$\therefore (9, 2)$  lies inside the circle.

M1 – realises the need to find distance from  $(9, 2)$  to  $C$

A1 – need to indicate  $< 5$  (radius) and therefore inside the circle

The line  $4y = 3x + 1$  meets the circle at the points  $P$  and  $T$ , and the  $x$ -axis at  $S$ .  
 $T$  lies between  $P$  and  $S$ .

(c) Without finding the coordinates of  $P$  and of  $T$ , find the length  $TS$ .

[4]

At  $S$ ,  $y = 0$

$$3x + 1 = 0$$

$$x = -\frac{1}{3}$$

$$S\left(-\frac{1}{3}, 0\right)$$

Subst  $x = 5$  into  $4y = 3x + 1$ ,

$$4y = 3(5) + 1$$

$$y = \frac{16}{4} = 4$$

$\therefore (5, 4)$  lies on the line  $4y = 3x + 1$

Distance from  $C$  to  $S$

$$= \sqrt{\left(5 - \left(-\frac{1}{3}\right)\right)^2 + 4^2}$$

$$= 6\frac{2}{3} \text{ units}$$

$$\therefore TS = 6\frac{2}{3} - 5$$

$$= 1\frac{2}{3} \text{ units}$$

M1 – realises that  $y = 0$  for  $S$

B1 – shows centre lies on line

M1 – distance from their  $C$  to  $S$

A1

- 11** The table shows, to 1 decimal place, the mass,  $m$  of a radioactive substance, in grams, after  $t$  days.

$t$	5	10	15	20	25
$m$	57.7	37.0	23.7	15.2	9.7

- (a) On the grid opposite, plot  $\ln m$  against  $t$  and draw a straight line graph. [2]
- (b) Find the gradient of your straight line and hence express  $m$  in the form  $m_0 e^{kt}$ , where  $m_0$  and  $k$  are constants. [4]

$$\text{gradient} = -\frac{2.25}{25} = -0.09$$

$$\ln m = -0.09t + 4.5$$

$$m = e^{-0.09t+4.5}$$

$$m = e^{-0.09t} \cdot e^{4.5}$$

$$m = 90.0e^{-0.09t}$$

B1 – finds gradient correctly

M1 – finds  $\ln m = 'm't + c$  with their gradient + y -intercept

A1,A1

The half-life of a radioactive substance is the length of time it takes for half of the substance to decay.

- (c) In order to determine the half-life of the radioactive substance, a suitable straight line can be drawn on the same set of axes as your graph. Find the equation of this line and hence determine the half-life of the radioactive substance. [3]

$$m = 90.0e^{-0.09t}$$

when  $t = 0$ ,

$m = 90.0$  (initial mass)

half of original mass = 45

$$\ln 45 = 3.80.$$

line to be drawn :

$$y = 3.8$$

Half-life = 7.75 days

M1 – attempt to find original mass

A1


A1



$t$	5	10	15	20	25
$\ln m$	4.055	3.61	3.16	2.72	2.27

P1 – at least 3 points

P2 – straight line drawn with all points plotted correctly



- 12** A particle travels in a straight line so that its velocity,  $v$  cm/s,  $t$  seconds after passing through a fixed point  $O$ , is given by  $v = t^2 - kt + 5$ , where  $k$  is a constant. The particle first comes to an instantaneous rest at the point  $P$  and then at the point  $Q$ .

(a) Given that the particle reaches a minimum velocity at  $t = 3$ , show that  $k = 6$ . [2]

$$v = t^2 - kt + 6$$

$$= \left(t - \frac{k}{2}\right)^2 + 5 - \frac{k^2}{4}$$

B1

minimum velocity occurs

$$\text{when } t - \frac{k}{2} = 0$$

$$3 - \frac{k}{2} = 0$$

√M1

$$6 - k = 0$$

$$k = 6 \text{ (shown)}$$

$$v = t^2 - kt + 6$$

$$\frac{dv}{dt} = 2t - k$$

$$\text{when } t = 3, \frac{dv}{dt} = 0$$

$$2(3) - k = 0$$

$$6 - k = 0$$

$$k = 6 \text{ (shown)}$$

B1

√M1 – recognises  $\frac{dv}{dt} = 0$  when  $t = 3$

AG

(b) Find the distance  $PQ$ . [6]

At  $P$  and  $Q$ ,  $v = 0$

$$t^2 - 6t + 5 = 0$$

$$(t-1)(t-5) = 0$$

$$t = 1 \text{ or } t = 5$$

$$s = \frac{1}{3}t^3 - 3t^2 + 5t + c$$

$$\text{when } t = 0, s = 0, \therefore c = 0$$

$$s = \frac{1}{3}t^3 - 3t^2 + 5t$$

$$\text{when } t = 1, s = 2\frac{1}{3}$$

$$\text{when } t = 5, s = -8\frac{1}{3}$$

$$\text{Distance } PQ = 2\frac{1}{3} + 8\frac{1}{3}$$

$$= 10\frac{2}{3} \text{ m}$$

M1 – equates  $v$  to 0

A1

M1 – realises the need to integrate

A1

M1 –  $s$  at their  $t = 1$  or  $t = 5$

A1

- (c) With working clearly shown, explain whether the particle will pass by  $O$  again, after the first 7 seconds.

[2]

$$\text{At } t = 7, s = 2\frac{1}{3}, v = 12$$

Since  $s > 0$ ,  $v > 0$  and there are no more turning points after  $t = 5$ , the particle will not return to  $O$  after 7 seconds.

**Alternative solution**

$$\frac{1}{3}t^3 - 3t^2 + 5t = 0$$

$$t^3 - 9t^2 + 15t = 0$$

$$t(t^2 - 9t + 15) = 0$$

$$t = 0 \text{ or } t = \frac{9 \pm \sqrt{(-9)^2 - 4(1)(15)}}{2}$$

$$= 6.79 \quad \text{or} \quad 2.21$$

Since the last time that the particle is at  $O$  is 6.79s, which is before 7 seconds, the particle will not pass by  $O$  again after 7 seconds.

M1 – finds  $s$  or  $v$  at  $t = 7$

A1 - explains no turning points  
particle will not return to  $O$

M1 – Attempt at solution for  
 $s = 0$

A1 – explains that last time  
that the particle is at  $O$  is  
6.79s , which is before 7s and  
hence does not return to  $O$

