

Name:	Class:	Class Register Number:
--------------	---------------	-------------------------------



中正中學

CHUNG CHENG HIGH SCHOOL (MAIN)

Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School
 Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School
 Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School
 Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School
 Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School Chung Cheng High School

**PRELIMINARY EXAMINATION 2023
SECONDARY 4**

ADDITIONAL MATHEMATICS

4049/02

Paper 2

Tuesday 29 August 2023

Candidates answer on the Question Paper.

2 hours 15 minutes

MARKS SCHEME

This document consists of **20** printed pages and **2** blank pages.

Mathematical Formulae**1. ALGEBRA***Quadratic Equation*

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1) \dots (n-r+1)}{r!}.$

2. TRIGONOMETRY*Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} ab \sin C$$

- 1 (a) The line $4x = 3y + 2$ intersects the curve $x^2 - xy + 5 = 0$ at the points A and B .
Find the midpoint of AB . [5]

$$4x = 3y + 2 \quad \dots(1)$$

$$x^2 - xy + 5 = 0 \quad \dots(2)$$

From (1):

$$3y = 4x - 2$$

$$y = \frac{4x - 2}{3} \quad \dots(3)$$

Sub. (3) into (2):

$$x^2 - x\left(\frac{4x - 2}{3}\right) + 5 = 0$$

$$3x^2 - 4x^2 + 2x + 15 = 0$$

$$-x^2 + 2x + 15 = 0$$

$$x^2 - 2x - 15 = 0$$

$$(x - 5)(x + 3) = 0$$

$$x = 5 \text{ or } -3 \text{ sub. into (3):}$$

$$y = 6 \text{ or } -\frac{14}{3}$$

$$\therefore A\left(-3, -\frac{14}{3}\right) \text{ and } B(5, 6)$$

$$\begin{aligned} \text{Midpoint of } AB &= \left(\frac{-3+5}{2}, \frac{-\frac{14}{3}+6}{2} \right) \\ &= \left(1, \frac{2}{3} \right) \end{aligned}$$

M1 f.t. – substitution

M1 f.t. – solving quadratic

M1 – either correct x or y values

M1 f.t. – midpoint formula

A1

- (b) Find the least value of the integer h for which $hx^2 + 5x + h$ is positive for all real values of x . [3]

For $hx^2 + 5x + h > 0$,

discriminant < 0

$$25 - 4(h)h < 0$$

$$25 - 4h^2 < 0$$

$$(5 - 2h)(5 + 2h) < 0$$

$$h < -\frac{5}{2} \quad \text{or} \quad h > \frac{5}{2}$$

Since $h > 0$, $h > \frac{5}{2}$.

\therefore Least integer value of $h = 3$.

B1 – discriminant

M1 f.t. – factorising quadratic

A1

- (c) Given that the line $y = 3x + p$ is tangent to the curve $y = x^2 + 5x + q$, where p and q are integers, prove that p and q are consecutive numbers. [4]

$$y = 3x + p \quad \dots (1)$$

$$y = x^2 + 5x + q \quad \dots (2)$$

Sub. (1) into (2):

$$3x + p = x^2 + 5x + q$$

$$x^2 + 5x + q - 3x - p = 0$$

$$x^2 + 2x + q - p = 0$$

$$a = 1, b = 2, c = q - p$$

Line is tangent to curve $\rightarrow b^2 - 4ac = 0$

$$2^2 - 4(1)(q - p) = 0$$

$$4 - 4q + 4p = 0$$

$$4q = 4 + 4p$$

$$q = 1 + p$$

Since $q = 1 + p$, q will always be the next number after p .

Hence, p and q are consecutive numbers (proved).

M1 – substitution

M1 – forming quadratic

M1 f.t. – any use of discriminant

A1 – with explanation

Alternative:

$$y = x^2 + 5x + q$$

$$\frac{dy}{dx} = 2x + 5$$

Since line is tangent to curve and gradient of line = 3,

$$2x + 5 = 3$$

$$2x = -2$$

$$x = -1$$

$$y = 3x + p \quad \dots (1)$$

$$y = x^2 + 5x + q \quad \dots (2)$$

Sub. (1) into (2) and $x = -1$:

$$3x + p = x^2 + 5x + q$$

$$3(-1) + p = (-1)^2 + 5(-1) + q$$

$$-3 + p = -4 + q$$

$$q - p = 1$$

Since the difference between q and p is 1, q will always be the next number after p .

Hence, p and q are consecutive numbers (proved).

M1 – equate $\frac{dy}{dx}$ to 3

M1 f.t. – finding x

M1 – substitution

A1 – with explanation

- 2 (a) By considering the general term in the binomial expansion of $\left(x^3 - \frac{2}{x}\right)^7$, explain why there are only odd powers of x in this expansion. [3]

$$\begin{aligned} T_{r+1} &= \binom{7}{r} (x^3)^{7-r} \left(-\frac{2}{x}\right)^r \\ &= \binom{7}{r} (x^{21-3r}) (-2)^r (x)^{-r} \\ &= \binom{7}{r} (-2)^r x^{21-4r} \end{aligned}$$

Power of $x = 21 - 4r$

Since **$4r$ is an even number** for all non-negative integer values of r and 21 is an odd number, then **$21 - 4r$ is always an odd number**.

Therefore, there are only odd powers of x in this expansion.

B1 – general formula

M1 f.t. – finding powers of x

A1 – conclusion

- (b) Find the term independent of x in the expansion of $\left(x^3 - \frac{2}{x}\right)^7 \left(\frac{5}{x} - 2x^2\right)$. [3]

$$\begin{aligned} \text{Consider } 21 - 4r &= 1, \\ 4r &= 20 \\ r &= 5 \end{aligned}$$

$$\begin{aligned} \left(x^3 - \frac{2}{x}\right)^7 \left(\frac{5}{x} - 2x^2\right) &= \left[\dots + \binom{7}{5} (-2)^5 x^{21-4(5)} + \dots \right] \left(\frac{5}{x} - 2x^2\right) \\ &= (\dots - 672x + \dots) \left(\frac{5}{x} - 2x^2\right) \end{aligned}$$

$$\begin{aligned} \text{Term independent of } x &= -672 \times 5 \\ &= -3360 \end{aligned}$$

M1 f.t. – identifying x term

M1 f.t. – expansion

A1

3 The expression $7 \sin \theta + 3 \cos \theta$ is defined for $0^\circ \leq \theta \leq 360^\circ$.

(a) Using $R \sin(\theta + \alpha)$, where $R > 0$ and $0^\circ < \alpha < 90^\circ$, solve the equation

$$7 \sin \theta = 5 - 3 \cos \theta.$$

[5]

$$\begin{aligned} 7 \sin \theta + 3 \cos \theta &= R \sin(\theta + \alpha) \\ &= R \sin \theta \cos \alpha + R \cos \theta \sin \alpha \end{aligned}$$

Comparing,

$$7 = R \cos \alpha \quad \text{and} \quad 3 = R \sin \alpha$$

$$\begin{aligned} R &= \sqrt{7^2 + 3^2} \\ &= \sqrt{58} \end{aligned}$$

$$\tan \alpha = \frac{3}{7}$$

$$\begin{aligned} \alpha &= \tan^{-1}\left(\frac{3}{7}\right) \\ &= 23.1985...^\circ \end{aligned}$$

$$\therefore 7 \sin \theta + 3 \cos \theta = \sqrt{58} \sin(\theta + 23.1985...^\circ)$$

M1 – attempt to find R

M1 – attempt to find α

[only accept $\tan^{-1}\left(\frac{3}{7}\right)$ or $\tan^{-1}\left(\frac{7}{3}\right)$]

$$7 \sin \theta = 5 - 3 \cos \theta$$

$$7 \sin \theta + 3 \cos \theta = 5$$

$$\sqrt{58} \sin(\theta + 23.1985...^\circ) = 5$$

$$\sin(\theta + 23.1985...^\circ) = \frac{5}{\sqrt{58}}$$

$$\text{basic angle} = \sin^{-1}\left(\frac{5}{\sqrt{58}}\right)$$

$$= 41.0359...^\circ$$

$$\theta + 23.1985...^\circ = 41.0359...^\circ \quad \text{or} \quad 180^\circ - 41.0359...^\circ$$

$$\theta = 17.8^\circ \quad \text{or} \quad 115.8^\circ \text{ (1 dec. pl.)}$$

M1 f.t. – substitute R -form

M1 f.t. – basic angle

A1

- (b) State the largest and smallest values of $(7 \sin \theta + 3 \cos \theta)^2 - 12$ and find the corresponding values of θ . [4]

$$\text{largest value of } (7 \sin \theta + 3 \cos \theta)^2 - 12 = (\sqrt{58})^2 - 12 \quad \text{or} \quad (-\sqrt{58})^2 - 12 \\ = 46$$

B1

$$\text{occurs when } \theta + 23.1985\dots^\circ = 90^\circ \quad \text{or} \quad 270^\circ \\ \theta = 66.8^\circ \quad \text{or} \quad 246.8^\circ \text{ (1 dec. pl.)}$$


B1

$$\text{smallest value of } (7 \sin \theta + 3 \cos \theta)^2 - 12 = (0)^2 - 12 \\ = -12$$

B1

$$\text{occurs when } \theta + 23.1985\dots^\circ = 180^\circ \text{ or } 360^\circ \\ \theta = 156.8^\circ \text{ or } 336.8^\circ \text{ (1 dec. pl.)}$$

B1



The diagram shows a circle passing through the points A, B, C, D and E . The straight line FDG is tangent to the circle at D while FAB and FEC are secant lines.

Given that angle $FDA = \text{angle } ADB$,

- (a) show that triangle ABD is an isosceles triangle,

[2]

$$\begin{aligned}\angle ABD &= \angle FDA && \text{(alternate segment theorem)} \\ &= \angle ADB\end{aligned}$$

B1

Since $\angle ABD = \angle ADB$, they form **base angles of isosceles triangle**.

B1

Thus, triangle ABD is an isosceles triangle. (shown)

- (b) prove that $AF \times BF = EF \times CF$.

[3]

$$\angle FAE = \angle FCB \quad \text{(exterior } \angle \text{ of cyclic quadrilateral)}$$

B1

$$\angle AFE = \angle CFB \quad \text{(common } \angle \text{)}$$


B1

Thus, triangle AFE is similar to triangle CFB .

$$\frac{AF}{CF} = \frac{EF}{BF} \quad \text{(ratio of corresponding sides are equal)}$$

B1

$$AF \times BF = EF \times CF \quad \text{(proved)}$$

**Alternative:**

$$\angle FBE = \angle FCA \quad (\angle\text{s in same segment})$$

$$\angle EFB = \angle AFC \quad (\text{common } \angle)$$

Thus, triangle EFB is similar to triangle AFC .

$$\frac{AF}{EF} = \frac{CF}{BF} \quad (\text{ratio of corresponding sides are equal})$$

$$AF \times BF = EF \times CF \quad (\text{proved})$$

B1

B1

B1

- 5 An object is heated in an oven until it reaches a temperature of X °C. It is then allowed to cool under room temperature. Its temperature, T °C, can be modelled by $T = 18 + 62e^{-kt}$, where t is the time in minutes since the object starts cooling.

- (a) Find the value of X .

[1]

$$\begin{aligned}\text{When } t = 0, \quad T &= X. \\ X &= 18 + 62e^{-k(0)} \\ &= 18 + 62 \\ &= 80\end{aligned}$$

B1

When $t = 10$, the temperature of the object is 65 °C.

- (b) Find the temperature of the object an hour later, giving your answer to one decimal place.

[5]

$$\begin{aligned}\text{When } t = 10, \quad T &= 65. \\ 65 &= 18 + 62e^{-k(10)} \\ \frac{47}{62} &= e^{-10k} \\ \ln \frac{47}{62} &= \ln e^{-10k} \\ \ln \frac{47}{62} &= -10k \\ k &= -\frac{1}{10} \ln \frac{47}{62} \\ &= 0.0276986...\end{aligned}$$

B1 – sub. 10 and 65 correctly

M1 – Take ln on both sides

M1 – finding k

$$\begin{aligned}\text{When } t = 60, \\ T &= 18 + 62e^{-\left(-\frac{1}{10} \ln \frac{47}{62}\right)(60)} \\ &= 29.765... \\ &= 29.8 \text{ °C (1 dec. pl.)}\end{aligned}$$

$$\begin{aligned}\text{When } t = 70, \\ T &= 18 + 62e^{-\left(-\frac{1}{10} \ln \frac{47}{62}\right)(70)} \\ &= 26.9193... \\ &= 26.9 \text{ °C (1 dec. pl.)}\end{aligned}$$

M1 f.t.– sub. $t = 60$ or 70

A1

- (c) What does the model suggest about the room temperature? Explain your answer.

[2]

Since $e^{-kt} > 0$, then $62e^{-kt} > 0$.
Thus, **as t becomes larger, $62e^{-kt}$ approaches zero.**
Therefore, T will **approach 18 °C** as t becomes larger.
Hence, the room temperature is **18 °C**.

B1 – explanation

B1 – temperature

- 6 The equation of a curve is $y = \ln\left(\frac{2x-1}{3x-1}\right)$, where $x > \frac{1}{2}$.

- (a) Find $\frac{dy}{dx}$, expressing it as a single fraction. [3]

$$\begin{aligned} y &= \ln\left(\frac{2x-1}{3x-1}\right) \\ &= \ln(2x-1) - \ln(3x-1) \end{aligned}$$

B1 – quotient law of logarithms

$$\begin{aligned} \frac{dy}{dx} &= \frac{2}{2x-1} - \frac{3}{3x-1} \\ &= \frac{2(3x-1) - 3(2x-1)}{(2x-1)(3x-1)} \\ &= \frac{1}{(2x-1)(3x+1)} \end{aligned}$$

M1 f.t. – differentiate either term

A1 – single fraction

Alternative:

$$\begin{aligned} \frac{dy}{dx} &= \frac{\frac{2(3x-1) - 3(2x-1)}{(3x-1)^2}}{\frac{2x-1}{3x-1}} \\ &= \frac{\frac{1}{(3x-1)^2}}{\frac{2x-1}{3x-1}} \\ &= \frac{1}{(2x-1)(3x+1)} \end{aligned}$$

B1 – differentiate \ln in the form $\frac{f'(x)}{f(x)}$

M1 – apply quotient rule to numerator.

A1 – single fraction

- (b) Explain why the curve will be almost parallel to the x -axis as x becomes very large. [2]

Since $(2x-1)(3x+1)$ becomes a very large number as x becomes very large, $\frac{dy}{dx}$ approaches **0**, the curve will **almost horizontal**, thus it will be almost parallel to the x -axis.

M1

A1

- (c) Find the value of x at the instant when the rate of change of x is twice the rate of change of y . [3]

$$\frac{dy}{dt} = \frac{dy}{dx} \times \frac{dx}{dt}$$

$$\frac{dy}{dt} = \frac{1}{(2x-1)(3x-1)} \times 2 \left(\frac{dy}{dt} \right)$$

$$1 = \frac{2}{(2x-1)(3x-1)}$$

$$(2x-1)(3x-1) = 2$$

$$6x^2 - 5x + 1 - 2 = 0$$

$$6x^2 - 5x - 1 = 0$$

$$(x-1)(6x+1) = 0$$

$$x = 1 \text{ or } -\frac{1}{6} \text{ (rej. } \because x > \frac{1}{2} \text{)}$$

M1 f.t. – substitution

B1 – substitute $\frac{dx}{dt} = 2 \frac{dy}{dt}$

A1

7 (a) Show that $\tan A + \cot A = 2\operatorname{cosec} 2A$.

[3]

Alternative:

$$\text{LHS} = \tan A + \cot A$$

$$= \tan A + \frac{1}{\tan A}$$

$$= \frac{\tan^2 A + 1}{\tan A}$$

$$= \frac{\sec^2 A}{\tan A}$$

$$= \frac{1}{\frac{\cos^2 A}{\sin A}}$$

$$= \frac{1}{\sin A \cos A}$$

$$= \frac{2}{2 \sin A \cos A}$$

$$= 2\operatorname{cosec} 2A \text{ (shown)}$$

$$\text{LHS} = \tan A + \cot A$$

$$= \frac{\sin A}{\cos A} + \frac{\cos A}{\sin A}$$

$$= \frac{\sin^2 A + \cos^2 A}{\sin A \cos A}$$

$$= \frac{1}{\sin A \cos A}$$

$$= \frac{2}{2 \sin A \cos A}$$

$$= \frac{2}{\sin 2A}$$

$$= 2\operatorname{cosec} 2A \text{ (shown)}$$

$$\text{M1} - \frac{\sin A}{\cos A} \text{ or } \frac{1}{\tan A}$$

$$\text{M1} - \text{for } \sin^2 A + \cos^2 A = 1 \\ \text{or for } \tan^2 A + 1 = \sec^2 A$$

A1 – award only if last 3 steps are shown

(b) Hence, solve the equation $\frac{1}{\tan A + \cot A} = \frac{1}{4}$ for $0 \leq A \leq 2\pi$.

[3]

$$\frac{1}{\tan A + \cot A} = \frac{1}{4}$$

$$\tan A + \cot A = 4$$

$$2\operatorname{cosec} 2A = 4$$

$$\operatorname{cosec} 2A = 2$$

$$\sin 2A = \frac{1}{2}$$

$$\text{basic } \angle = \sin^{-1}\left(\frac{1}{2}\right)$$

$$= \frac{\pi}{6}$$

$$2A = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{13\pi}{6}, \frac{17\pi}{6}$$

$$A = \frac{\pi}{12}, \frac{5\pi}{12}, \frac{13\pi}{12}, \frac{17\pi}{12}$$

B1 – substitution of (a)


$$\text{M1} - \text{for finding } \frac{\pi}{6} \text{ and } \frac{5\pi}{6}$$

A1

- (c) The diagram shows, for $0 \leq x \leq \pi$, the curve $y = \sin 2x$ and the line $y = \frac{1}{2}$.

Showing all your working, find the area of the shaded region.

[5]



Area bounded by curve, x -axis and lines $x = 0$ and $x = \frac{\pi}{12}$

$$\begin{aligned}
 &= \int_0^{\frac{\pi}{12}} \sin 2x \, dx \\
 &= \left[\frac{-\cos 2x}{2} \right]_0^{\frac{\pi}{12}} \\
 &= \left[-\frac{\cos \frac{\pi}{6}}{2} \right] - \left[-\frac{\cos 0}{2} \right] \\
 &= \left(-\frac{\sqrt{3}}{4} + \frac{1}{2} \right) \text{ units}^2 \quad \text{or} \quad 0.066987... \text{ units}^2
 \end{aligned}$$

M1 – integrate $\sin 2x$

DM1 – evaluate with either
limits 0 to $\frac{\pi}{12}$ or $\frac{5\pi}{12}$ to $\frac{\pi}{2}$ or
 $\frac{\pi}{2}$ to $\frac{7\pi}{12}$


$$\begin{aligned}
 \text{Area of rectangle} &= \left(\frac{5\pi}{12} - \frac{\pi}{12} \right) \times \frac{1}{2} \\
 &= \frac{\pi}{6} \text{ units}^2
 \end{aligned}$$

M1 – area of rectangle

$$\begin{aligned}
 \text{Area of shaded region} &= 3 \times \left(-\frac{\sqrt{3}}{4} + \frac{1}{2} \right) + \frac{\pi}{6} \\
 &= 0.725 \text{ units}^2 \quad (3 \text{ sig.fig})
 \end{aligned}$$

M1 f.t. – area under curve $\times 3$

A1



The diagram shows a parallelogram with vertices A , $B(6, 21)$, C and $D(3, 0)$. The point $E(8, 17)$ lies on BC . The line CD makes an angle θ with the positive x -axis such that $\tan \theta = 1$. A line is drawn, parallel to the y -axis, from A to meet the x -axis at N .

(a) Show that the coordinates of A are $(-3, 12)$.

[5]

$$\begin{aligned} m_{AB} &= m_{CD} \\ &= 1 \end{aligned}$$

B1 – gradient = $\tan \theta$

$$\begin{aligned} \text{Eq. of } AB: \quad y - 21 &= (1)(x - 6) \\ y &= x + 15 \quad \dots(1) \end{aligned}$$

$$\begin{aligned} m_{AD} &= m_{BE} \\ &= \frac{17 - 21}{8 - 6} \\ &= -2 \end{aligned}$$

B1 – gradient formula

$$\begin{aligned} \text{Eq. of } AD: \quad y - 0 &= -2(x - 3) \\ y &= -2x + 6 \quad \dots(2) \end{aligned}$$

M1 f.t. – finding equation of AD or AB

Sub. (1) into (2):

$$x + 15 = -2x + 6$$

$$3x = -9$$

$$x = -3 \text{ sub. into (1):}$$

$$y = 12$$

$\therefore A(-3, 12)$ (shown)

M1 f.t. – substitution

A1 (A.G.)

Alternative:

$$\begin{aligned}\text{Eq. of } CD: \quad y - 0 &= (1)(x - 3) \\ y &= x - 3 \quad \dots(1)\end{aligned}$$

$$\begin{aligned}m_{BC} &= m_{BE} \\ &= \frac{17 - 21}{8 - 6} \\ &= -2\end{aligned}$$

$$\begin{aligned}\text{Eq. of } BC: \quad y - 21 &= -2(x - 6) \\ y &= -2x + 33 \quad \dots(2)\end{aligned}$$

Sub. (1) into (2):

$$x - 3 = -2x + 33$$

$$3x = 36$$

$$x = 12 \text{ sub. into (1):}$$

$$y = 9$$

$$\therefore C(12, 9)$$

Midpoint of AC = Midpoint of BD

$$\left(\frac{x+12}{2}, \frac{y+9}{2} \right) = \left(\frac{3+6}{2}, \frac{0+21}{2} \right)$$

$$\left(\frac{x+12}{2}, \frac{y+9}{2} \right) = \left(\frac{9}{2}, \frac{21}{2} \right)$$

Comparing,

$$\frac{x+12}{2} = \frac{9}{2} \quad \frac{y+9}{2} = \frac{21}{2}$$

$$x+12 = 9 \quad \text{and} \quad y+9 = 21$$

$$x = -3 \quad y = 12$$

$$\therefore A(-3, 12) \text{ (shown)}$$

B1 – gradient formula

M1 f.t. – finding equation of BC

M1 f.t. – substitution

M1 f.t. – equating midpoints

A1 (A.G.)

(b) Hence, find the area of parallelogram $ABCD$.

[2]

area of parallelogram $ABCD = 2 \times$ area of triangle ABD

$$\begin{aligned}
 &= 2 \times \frac{1}{2} \begin{vmatrix} -3 & 3 & 6 & -3 \\ 12 & 0 & 21 & 12 \end{vmatrix} \\
 &= 2 \times \frac{1}{2} [(63 + 72) - (-63 + 36)] \\
 &= 135 + 27 \\
 &= 162 \text{ units}^2
 \end{aligned}$$

M1 – allow clockwise direction

A1

Alternative:

If students have found the coordinates of C in part (a),

$$\begin{aligned}
 \text{area of parallelogram } ABCD &= \frac{1}{2} \begin{vmatrix} 3 & 12 & 6 & -3 & 3 \\ 0 & 9 & 21 & 12 & 0 \end{vmatrix} \\
 &= \frac{1}{2} [(27 + 252 + 72) - (36 - 63 + 54)] \\
 &= \frac{1}{2} (351 - 27) \\
 &= 162 \text{ units}^2
 \end{aligned}$$

M1 f.t. – allow clockwise direction

A1

- (c) A point F with y -coordinate of 5 lies on the line CD .
Explain why $AEFN$ is a parallelogram.

[5]

$$\begin{aligned}\text{Eq. of } CD: \quad y - 0 &= (1)(x - 3) \\ y &= x - 3 \quad \dots(3)\end{aligned}$$

Let coordinates of F be $(x, 5)$.

Sub. $(8, y)$ into (3) :

$$\begin{aligned}y &= 8 - 3 \\ &= 5\end{aligned}$$

$\therefore F(8, 5)$

Coordinates of $N = (-3, 0)$

$$\begin{aligned}\text{length of } AN &= \sqrt{(-3+3)^2 + (12-0)^2} \\ &= 12 \text{ units}\end{aligned}$$

$$\begin{aligned}\text{length of } EF &= \sqrt{(8-8)^2 + (17-5)^2} \\ &= 12 \text{ units}\end{aligned}$$

Since F lies directly below E , the line EF is a vertical line.
Thus, the lines EF and AN are parallel.

And that the lengths of AN and EF are equal (i.e., 12 units),

they form a pair of parallel and equal opposite sides.
Thus, $AEFN$ is a parallelogram.

Alternative 1:

$$\begin{aligned}\text{gradient of } AE &= \frac{17-12}{8-(-3)} & \text{gradient of } FN &= \frac{5-0}{8-(-3)} \\ &= \frac{5}{11} & &= \frac{5}{11}\end{aligned}$$

Since gradients of lines AE and FN are the same, the lines AE and FN are parallel.

Since F lies directly below E , the line EF is a vertical line.
Thus, the lines EF and AN are parallel.

Since there are 2 pairs of parallel lines, $AEFN$ is a parallelogram.

B1

M1 f.t. – substitution

M1 f.t. – finding either lengths

A1 – vertical lines are parallel

A1 – conclusion

M1 f.t. – finding either gradients

A1 – vertical lines are parallel

A1 – conclusion

Alternative 2:

$$\begin{aligned}\text{Midpoint of } AF &= \left(\frac{-3+8}{2}, \frac{12+5}{2} \right) \\ &= \left(\frac{5}{2}, \frac{17}{2} \right)\end{aligned}$$

M1 f.t. – midpoint of AF

$$\begin{aligned}\text{Midpoint of } NE &= \left(\frac{-3+8}{2}, \frac{0+17}{2} \right) \\ &= \left(\frac{5}{2}, \frac{17}{2} \right)\end{aligned}$$

M1 f.t. – midpoint of AE

Since the diagonals intersect at the same point, $AEFN$ is a parallelogram.

A1 – conclusion

- 9 (a) Differentiate $x \ln x$ with respect to x .

[2]

$$\begin{aligned}\frac{d}{dx}(x \ln x) &= x \left(\frac{1}{x} \right) + (1) \ln x \\ &= 1 + \ln x\end{aligned}$$

B1, B1 – for each term

- (b) A curve $y = f(x)$ is such that $\frac{d^2y}{dx^2} = 24x^2 + \frac{16}{x}$, where $x > 0$. The line $y = 24x - 40$ is parallel to the tangent of the curve at $P(1, -16)$.

By using the result found in part (a), find the equation of the curve.

[6]

$$\begin{aligned}\frac{dy}{dx} &= \int 24x^2 + \frac{16}{x} dx \\ &= \frac{24x^3}{3} + 16 \ln x + c \\ &= 8x^3 + 16 \ln x + c, \text{ where } c \text{ is a constant.}\end{aligned}$$

B1, B1 – for each integral

$$\begin{aligned}\text{When } x = 1, \frac{dy}{dx} &= 24. \\ 24 &= 8(1)^3 + 16 \ln(1) + c \\ c &= 16\end{aligned}$$

M1 f.t. – finding c

$$\therefore \frac{dy}{dx} = 8x^3 + 16 \ln x + 16$$

$$\begin{aligned}y &= \int 8x^3 + 16 \ln x + 16 dx \text{ or } y = \int 8x^3 + 16(1 + \ln x) dx \\ &= \frac{8x^4}{4} + 16 \left(x \ln x - \int 1 dx \right) + 16x + c_1 \\ &= \frac{8x^4}{4} + 16(x \ln x - x + c_2) + 16x + c_1 \\ &= 2x^4 + 16x \ln x - 16x + 16x + c_3 \\ &= 2x^4 + 16x \ln x + c_3, \text{ where } c_1, c_2 \text{ and } c_3 \text{ are constants.}\end{aligned}$$

M1 f.t. – reverse

$$\begin{aligned}\text{When } x = 1, y &= -16. \\ -16 &= 2(1)^4 + 16(1) \ln(1) + c_3 \\ -16 &= 2 + c_3 \\ c_3 &= -18\end{aligned}$$

M1 f.t. – finding c_3

$$\therefore y = 2x^4 + 16x \ln x - 18$$

A1

Continuation of working space for question 9(b).

- (c) Explain why the condition $x > 0$ is necessary.


[1]

$\ln x$ is defined for $x > 0$.

or

$\ln x$ is undefined for $x < 0$.

B1



The diagram shows a solid prism with right-angled triangular ends that are perpendicular to the parallel sides AD , BE and CF , which are each y cm in length. The right-angled triangular ends have sides AC and DF , which are $3x$ cm, and sides BC and EF , which are $4x$ cm.

Given that the volume of the prism is 1200 cm^3 ,

- (a) find an expression for y in terms of x ,

[2]

$$\text{Vol. of prism} = \frac{1}{2}(3x)(4x)y$$

$$1200 = 6x^2y$$

$$y = \frac{200}{x^2}$$

B1 – volume formula

B1

- (b) show that the total surface area of the prism, $S \text{ cm}^2$, is given by $S = 12x^2 + \frac{2400}{x}$. [3]

$$\begin{aligned} AB &= \sqrt{(3x)^2 + (4x)^2} \\ &= \sqrt{25x^2} \\ &= 5x \text{ cm} \end{aligned}$$

B1 – finding length of AB

$$\begin{aligned} \text{Total surface area} &= 2 \times \frac{1}{2} (3x)(4x) + 3xy + 4xy + 5xy \\ &= 12x^2 + 12xy \\ &= 12x^2 + 12x \left(\frac{200}{x} \right) \\ &= 12x^2 + \frac{2400}{x} \text{ (shown)} \end{aligned}$$

M1 – at least 3 correct surfaces

A1 (A.G.)

- (c) Given that x can vary, find the value of x for which the total surface area of the prism is a stationary value. [3]

$$\frac{dS}{dx} = 24x - \frac{2400}{x^2}$$

B1 – differentiation

$$\text{For } S \text{ to be stationary, } \frac{dS}{dx} = 0$$

$$24x - \frac{2400}{x^2} = 0$$

M1 f.t.

$$24x = \frac{2400}{x^2}$$

$$x^3 = 100$$

$$x = \sqrt[3]{100}$$

$$= 4.64 \text{ (3 sig. fig.)}$$

A1

(d) Explain why this value of x gives the smallest surface area possible.

[2]

$$\frac{d^2S}{dx^2} = 24 + \frac{4800}{x^3}$$

When $x = \sqrt[3]{100}$,

$$\frac{d^2S}{dx^2} = 24 + \frac{4800}{\left(\sqrt[3]{100}\right)^3}$$

$$= 72 > 0 \quad \therefore \text{minimum}$$

Since $\frac{d^2S}{dx^2} > 0$, the surface area is the smallest when $x = 4.64$.

M1 f.t. – differentiation or
1st derivative test

A1 – conclude ‘> 0’ (must
show ‘72’)

BLANK PAGE