



**HILLGROVE SECONDARY SCHOOL
PRELIMINARY EXAMINATION 2023
SECONDARY FOUR (EXPRESS)**

CANDIDATE
NAME

()

CLASS

	-	
--	---	--

CENTRE
NUMBER

S				
---	--	--	--	--

INDEX
NUMBER

--	--	--	--

Additional Mathematics

4049/01

Paper 1

23 August 2023

Candidates answer on the Question Paper.

2 hours 15 minutes

No Additional Materials are required.

10.45 a.m. – 1.00 p.m.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, index number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an approved scientific calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 90.

Parent's/ Guardian's Signature: _____

Setter: Mdm Lee Li Lian

For Examiner's Use	
TOTAL	90

This document consists of **18** printed pages, including this page.

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} ab \sin C$$

1 The equation of a curve is $y = 4x^2 - 16x + 19$.

(a) By expressing $4x^2 - 16x + 19$ in the form $a(x+b)^2 + c$, where a , b and c are constants, [2]
find the coordinates of the stationary point on the curve.

(b) The line $y = 4x + 3$ intersects the curve at the points A and B . Find the value of k for [4]
which the distance AB can be expressed as $3\sqrt{k}$.

- 2 It is given that $(\sqrt{5})^9 - (\sqrt{5})^7 + (\sqrt{5})^5 - (\sqrt{5})^3 + 105(\sqrt{5}) = 5^k$. By factorisation, find the value of k . [4]

3

The loudness of a sound can be measured using the equation $L = 10 \lg \frac{I}{I_0}$, where I is the intensity of sound to be measured and I_0 is the intensity of sound that can barely be heard, also known as the threshold of hearing. The unit of L is the decibel (dB).

- (a) Given that the loudness of a scream is 110 dB, find the ratio of the intensity of the scream to the threshold of hearing. [2]

- (b) 130 dB is the pain threshold (the maximum level of sound we can hear without feeling intense pain and instantly damaging our hearing). [2]

Explain the impact, on hearing, the loudness of the sound of an unknown object falling from the sky onto Earth, if it has a sound intensity of $10^{-10.5}$ units and threshold of hearing of 10^{-25} units.

- 4 Given that the range of values of x where $x^2 + ax < b$ is $-4 < x < 5$, find the value of a and of b . [4]

5 The line $y = mx + c$ is drawn on the same axes as the curve $y = 4x - 2x^2$.

(a) Given that the line is a tangent to the curve when $c = \frac{1}{2}$, find the possible values of m . [3]

(b) The line $y = mx + c$ has a negative y -intercept. Do the line and the curve have 0, 1 or 2 points of intersections? Show your working clearly. [3]

6 A polynomial, P , is $3x^3 + 5x^2 - x + k$, where k is a constant.

(a) Find the value of k given that P leaves a remainder of 24 when divided by $x - 2$. [2]

(b) In the case where $k = -3$, the quadratic expression $3x^2 + ax^2 - 3$ is a factor of P . Find the value of the constant a . [4]

7 (a) Divide $2x^3 + 4x - 2$ by $x^3 + 2x$. [1]

(b) Express $\frac{2x^3 + 4x - 2}{x^3 + 2x}$ in partial fractions. [5]

(c) Hence, find $\int \frac{2x^3 + 4x^2 - 2}{x^3 + 2x} dx$. [2]

8 The line $3y + 4x = 12$ meets the x -axis at the point A and the y -axis at the point B .

(a) Write the coordinates of A and of B .

[2]

The perpendicular bisector of AB meets the line $y = x$ at the point C .

(b) Find the coordinates of C .

[4]

- (c) Find the area of quadrilateral $OACB$, where O is the origin.

[2]

- 9 (a)** Write down the first three terms in the expansion of $\left(2 - \frac{x}{4}\right)^n$, where n is a positive integer greater than 2, in ascending powers of x . [4]

The first two non-zero terms in the expansion of $(2 + x)\left(2 - \frac{x}{4}\right)^n$ in ascending powers of x are $a + bx^2$, where a and b are constants.

- (b)** Find the value of n . [2]

- (c)** Hence, find the value of a and of b . [2]

- 10** A curve is such that $\frac{d^2y}{dx^2} = 3e^{-x} + 8e^{2x}$. The curve intersects the y-axis at $P(0, -5)$ and has a gradient of 5 at P . Find the equation of the curve. [7]

- 11 (a)** Show that $6\sin^2 x - 4\cos^2 x$ can be written as $a + b \cos 2x$, where a and b are integers. [2]

Hence,

- (b)** state the period and amplitude of $6\sin^2 x - 4\cos^2 x$, [2]

- (c)** Sketch the graph of $y = 6\sin^2 x - 4\cos^2 x$ for $0 \leq x \leq 2\pi$ radians. [3]

- 12** Liquid is poured, at a constant rate of $25\pi \text{ cm}^3/\text{s}$, into a hemispherical bowl of radius $r \text{ cm}$.


When the depth of the liquid directly below the centre of the bowl is $x \text{ cm}$, the volume, $V \text{ cm}^3$, of the liquid in the bowl is given by $V = \frac{1}{3}\pi x^2(3r - x)$.

It is given that the radius of hemispherical bowl of radius is 12 cm , find

- (a) the time taken for the depth of the liquid directly below the centre of the bowl to reach 6 cm , [3]
- (b) the rate of change of the depth of liquid directly below the centre of the bowl at this time. [4]

13 (a) Show that $4\sec\theta + \tan\theta = 3\cot\theta$ can be expressed as $4\sin^2\theta + 4\sin\theta - 3 = 0$. [3]

(b) Hence, solve $4\sec 2x + \tan 2x = 3\cot 2x$ for $-180^\circ < x < 180^\circ$. [5]



The diagram shows triangles ABC and BCD whose vertices lie on the circumference of a circle. The chords BD and AC intersect at E and AC is parallel to FG . FG is a tangent to the circle at B .

Prove that

(a) $\triangle BCD$ is similar to $\triangle BEC$, [3]

(b) $BC^2 = BD \times BE$, [2]

(c) $\triangle ABC$ is an isosceles triangle.

[2]

Answer Key

- 1 (a) $(2, 3)$
(b) $k = 17$
- 2 $k = 4\frac{1}{2}$
- 3 (a) $10^{11} : 1$
- 4 $a = -1$ and $b = 20$
- 5 (a) $m = 6$ or $m = 2$
- 6 (a) $k = -18$
(b) $a = 2$
- 7 (a) $\frac{2x^3 + 4x - 2}{x^3 + 2x} = 2 - \frac{2}{x^3 + 2x}$
(b) $\frac{2x^3 + 4x - 2}{x^3 + 2x} = 2 - \frac{1}{x} + \frac{x}{x^2 + 2}$
(c) $2x - \ln x + \frac{1}{2} \ln(x^2 + 2) + c$ where c is a constant.
- 8 (a) $A = (3, 0)$ and $B = (0, 4)$
(b) $C = \left(3\frac{1}{2}, 3\frac{1}{2}\right)$
(c) 12.25 units^2
- 9 (a) $2^n - n(2^{n-3})x + n(n-1)2^{n-7}x^2 + \dots$
(b) $n = 4$
(c) $a = 32$ and $b = -5$
- 10 $y = 3e^{-x} + 2e^{2x} + 4x - 10$
- 11 (a) $1 - 5 \cos 2x$
(b) Period $= \pi$
Amplitude $= 5$
- 12 (a) 14.4 seconds
(b) $\frac{dx}{dt} = \frac{25}{108} \text{ cm/s}$
- 13 (b) $x = -165^\circ, -105^\circ, \underline{15^\circ, 75^\circ}$