



**TANJONG KATONG GIRLS' SCHOOL  
PRELIMINARY EXAMINATION  
SECONDARY FOUR EXPRESS**

CANDIDATE  
NAME

--

CLASS

	E	
--	---	--

INDEX  
NUMBER

--	--

---

**ADDITIONAL MATHEMATICS**

**4049/02**

PAPER 2

**16 August 2023**

**2 hour 15 minutes**

Candidates answer on the Question Paper

---

**READ THESE INSTRUCTIONS FIRST**

Write your index number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

Answer **all** questions.

If working is needed for any question it must be shown with the answer.

Omission of essential working will result in loss of marks.

The use of an approved scientific calculator is expected, where appropriate.

If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place.

For  $\pi$ , use either your calculator value or 3.142, unless the question requires the answer in terms of  $\pi$ .

The number of marks is given in brackets [ ] at the end of each question or part question.

The total marks for this paper is 90.

**For Examiner's use**

--

## 1. ALGEBRA

### *Quadratic Equation*

For the equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

### *Binomial expansion*

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

## 2. TRIGONOMETRY

### *Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$


### *Formula for $\Delta ABC$*

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

- 1 The diagram below shows the curve  $y = 4e^{2x} - 7$  and the curve  $y = 2e^{-2x}$ .



- (a) Using the diagram above, determine with explanation, the number of solutions for the equation  $4e^{2x} - 2e^{-2x} = 7$ . [1]

- (b) Solve the equation  $4e^{2x} - 7 = 2e^{-2x}$ . [4]


- 2 (a) Show that 1 is a solution of the equation  $x^3 - 5x^2 = 2 - 6x$ . Hence solve the equation  $x^3 - 5x^2 = 2 - 6x$  completely, expressing non-integer roots in surd form. [4]

- (b) Hence solve the equation  $(x-1)^3 - 5(x-1)^2 + 6x - 8 = 0$  completely.  
Express non-integer roots in surd form.

[2]

3      (a)    Given that  $y = \frac{4x}{\sqrt{3-2x}}$ , show that  $\frac{dy}{dx} = \frac{12-4x}{\sqrt{(3-2x)^3}}$ . [4]

(b) Hence find the value of  $\int_0^1 \frac{3-x}{\sqrt{(3-2x)^3}} dx$ . [4]



In the diagram,  $BE$  intersects  $AD$  at  $F$ .  $AF = 7$  cm and  $FD = 4$  cm.  $\angle ABF$ ,  $\angle ACD$  and  $\angle DEF$  are right angles.  $BCDF$  is a trapezium and  $\theta$  is an acute angle.

- (a) Show that the perimeter of  $BCDF$ ,  $P$  cm, is given by

$$P = 4 + 18\sin \theta + 4\cos \theta.$$

[2]

- (b) Express  $P$  in the form  $P = a + R\sin(\theta + \alpha)$ , where  $a$  is a constant,  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ .

[2]



(c) Find the maximum exact value of  $P$  and the corresponding value of  $\theta$ . [3]

(d) Find the value of  $\theta$  when  $P = 11$ . [2]

(e) Without evaluating  $\theta$ , determine whether the perimeter of  $BCDF$  can have a value of 28 cm. Justify your answer. [1]

- 5 (a) Find the set of values of  $x$  for which the curve  $y = -x^2 + 2x + 4$  lies below another curve  $y = x^2 - 2x - 2$  and represent this set on a number line. [4]

- (b) The line  $y = 2x + k$  is a normal to the curve  $y = -x^2 + 2x + 4$  at the point A.

- (i) Find the  $x$ -coordinate of A. [3]

- (ii) Find the value of the constant  $k$ . [2]

- 6 (a) (i) Write down, and simplify, the first 3 terms in the expansion of  $(2-x)^7$  in ascending powers of  $x$ . [2]

- (ii) Find the coefficient of  $x^2$  in the expansion of  $(1-8x+24x^2)(2-x)^7$ . [3]


- (b) (i) Using the general term, find the term in  $\frac{1}{x^7}$  in the binomial expansion of  $\left(2x - \frac{1}{x^2}\right)^{17}$ . [3]

- (ii) Explain why there is no term in  $\frac{1}{x^2}$  in the expansion of  $\left(2x - \frac{1}{x^2}\right)^{17}$ . [2]

- 7 (a) Designer bag  $X$  was first released in 1950. The price,  $\$V$ , of the bag is related to  $t$ , the number of years since 1950, by the formula  $V = ae^{kt}$ , where  $a$  and  $k$  are constants. The table below gives the value of bag  $X$  in 1984, 2002, 2008, and 2012.

Year	1984	2002	2008	2012
$t$ (years)	34	52	58	62
$V$ (\$)	1150	2850	4000	4900

- (i) On graph paper, plot  $\ln V$  against  $t$  and draw a straight line graph.  
 Use a scale of 2 cm to 0.5 on the vertical  $\ln V$ -axis, starting from  $\ln V = 5.0$ .  
 Use a scale of 2 cm to 10 years on the  $t$ -axis, starting from  $t = 0$ . [2]



(ii) Use your graph to estimate the value of  $a$  and of  $k$ . [3]

(iii) Estimate the year that the value of the bag will hit \$7000. [2]

(b) The variables  $x$  and  $y$  are related by the equation  $y = \frac{a}{x-b}$  where  $a$  and  $b$  are constants. Express the equation in a form suitable for drawing a straight line graph, and explain how the values of  $a$  and  $b$  may be obtained from the graph. [4]

- 8** A motorist, travelling at a constant velocity of  $V$  m/s, passed a fixed point  $X$  and saw few vehicles ahead. He stepped on the accelerator and his subsequent velocity,  $v$  m/s, is given by  $v = 60e^{\frac{t}{6}}$ , where  $t$  is the time in seconds after passing  $X$ . As he passed a point  $Y$ , his velocity has increased to twice his velocity at  $X$ .

**(a)** Find the time taken to travel from  $X$  to  $Y$ . [3]

**(b)** Find the acceleration of the motorist as he passes  $Y$ . [2]



(c) Find the distance  $XY$ .


[3]

- 9 (a) A circle,  $C_1$ , has the equation  $x^2 + y^2 + 2x - 6y = 10$ .

Find the coordinates of the centre and the exact radius of the circle.

[3]

(b)



$A$ ,  $B$ ,  $E$  and  $F$  are 4 points on another circle,  $C_2$ .

$EF$  is the perpendicular bisector of chord  $AB$ .  $EF$  cuts  $AB$  at the point  $\left(-\frac{9}{2}, \frac{11}{2}\right)$ .

The coordinates of point  $A$  are  $(-2, 8)$ .

- (i) Find the coordinates of  $B$ . [3]


- (ii) Explain why angle  $EAF = 90^\circ$ . [1]

The equation of the perpendicular bisector of chord  $AB$  is  $y + x = 1$ .  
A line  $2y = x + 8$  also passes through the centre of the circle.

- (iii) Find the centre of the circle. [2]

$P(2, 0)$  is another point on the circle.

- (iv) Find the equation of the tangent to the circle at  $P$ . [3]




The diagram shows the curve  $y = 4 - 2\cos 2x$  for  $0 \leq x \leq \frac{3}{2}\pi$  radians and a straight line passing through points  $P$  and  $R$ . The coordinates of  $P$  are  $(0, -1)$  and  $R$  is a minimum point of the curve.


Find the area of the region bounded by the curve  $y = 4 - 2\cos 2x$ , the line segment  $PR$  and the  $y$ -axis. [11]

Continuation of working space for question **10**.

**BLANK PAGE**



Question No.	Answer
1(a)	1
1(b)	0.347
2(a)	$1, 2 \pm \sqrt{2}$
2(b)	$2, 3 + \sqrt{2}$ or $3 - \sqrt{2}$
3(a)	Proof
3(b)	1
4(a)	Proof
4(b)	$P = 4 + 2\sqrt{85} \sin(\theta + 12.5^\circ)$
4(c)	$4 + 2\sqrt{85}, 77.5^\circ$
4(d)	$9.8^\circ$
4(e)	$P$ cannot a value of 18 cm.
5(a)	$\{x : x \in \mathbb{R}, x < -1 \text{ or } x > 3\}$ 
5(b)(i)	$\frac{5}{4}$
5(b)(ii)	$\frac{39}{16}$
6a(i)	$128 - 448x + 672x^2 + \dots$
6a(ii)	7328
6b(i)	$\frac{12446720}{x^7}$
6b(ii)	$r = \frac{19}{3}$ is not a non-negative integer in range $0 \leq r \leq 17$ . $\therefore$ there is no term in $\frac{1}{x^2}$ .

Question No.	Answer
7(a)(i)	
7(a)(ii)	$a = 200.3, k = 0.0518$
7(a)(iii)	2018
7(b)	$yx = a + by$ Plot $yx$ against $y$ to obtain a straight line graph with $a$ as the intercept on the vertical axis ( $yx$ -intercept) and $b$ as the gradient of the line. Thus, $a$ and $b$ may be obtained.
8(a)	4.16 s
8(b)	$20 \text{ m/s}^2$
8(c)	360 m
9(a)	$(-1, 3), 2\sqrt{5}$ units
9(b)(i)	$(-7, 3)$
9b(iii)	$(-2, 3)$
9b(ii)	Since EF is $\perp$ bisector of chord AB, $\therefore$ EF passes through centre of the circle (symmetrical property of circle). $\therefore$ EF is a diameter of the circle. Hence angle EAF = $90^\circ$ ( $\angle$ in a semicircle)
9b(iv)	$3y = 4x - 8$
10	11.0 square units